ASYMPTOTIC EQUILIBRIUM
OF ORDINARY DIFFERENTIAL SYSTEMS
IN A BANACH SPACE

by

A. Richard Mitchell and Roger W. Mitchell

Technical Report No. 7

March 1974

A differential system

$$x' = f(t, x) \quad \text{where } f \in C[[t_0, \infty) \times B, B] \quad (1)$$

$$x(\hat{t}) = \hat{x} \quad (2)$$

has asymptotic equilibrium if 1) for any initial condition $(\hat{t}, \hat{x}) \in [t_0, \infty) \times B$ the system has a solution $x(t)$ existing on $[\hat{t}, \infty)$ and such that $\lim_{t \rightarrow \infty} x(t)$ exists and is finite, and 2) for any $v \in B$ there exists $(\hat{t}, \hat{x}) \in [t_0, \infty) \times B$ and a solution $x(t)$ of (1)-(2) with $\lim_{t \rightarrow \infty} x(t) = v$.

Several papers have appeared dealing with asymptotic equilibrium of (1)-(2) when $B = \mathbb{R}^n$, and f is majorized by a scalar function $g(t, u)$ which is monotone in u for each t , [1,3]. However, when B is an arbitrary Banach space additional restrictions must be placed on f , (see [p.161,4; 5]). In [7] a set of sufficient conditions for local existence of solutions of (1)-(2) in an arbitrary Banach space is given. These conditions include the use of the Kuratowski measure of non-compactness of bounded sets, denoted throughout this paper by α (see [2,7]). Since our goal will be to give sufficient conditions for asymptotic equilibrium of (1)-(2) using α , the first lemma incorporates some known properties of α (see [7]).

Lemma 1. Let A and B be bounded subsets of a Banach space and $\{a_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$ be bounded sequences in the Banach space.

Then

$$(1.1) \quad \alpha(A + B) \leq \alpha(A) + \alpha(B) \quad \text{where } A + B = \{a + b \mid a \in A \text{ and } b \in B\}.$$

$$(1.2) \quad \alpha(\lambda A) = |\lambda| \alpha(A) \quad \text{for any scalar } \lambda.$$

$$(1.3) \quad \alpha\left\{\{a_n\}_{n=1}^{\infty}\right\} - \alpha\left\{\{b_n\}_{n=1}^{\infty}\right\} \leq \alpha\left\{\{a_n - b_n\}_{n=1}^{\infty}\right\}.$$

$$(1.4) \quad \alpha(A) = 0 \text{ iff } \bar{A} \text{ is compact (i.e. } \alpha(A) = 0 \text{ iff } A \text{ is relatively compact).}$$

$$(1.5) \quad \text{If } \|a\| \leq \epsilon \text{ for all } a \in A, \text{ then } \alpha(A) \leq 2\epsilon.$$

The proof of the following theorem may be found in [5].

Theorem 2. If B is a Banach space and

$$(2.1) \quad (1)-(2) \text{ has local solutions for all } (\hat{t}, \hat{x}) \in [t_0, \infty) \times B;$$

$$(2.2) \quad \|f(t, x)\| \leq g(t, \|x\|) \text{ for all } (t, x) \in [t_0, \infty) \times B;$$

$$(2.3) \quad g \in C[[t_0, \infty) \times \mathbb{R}^+, \mathbb{R}^+] \text{ is non-decreasing in } u \geq 0 \text{ for each } t \in [t_0, \infty);$$

$$(2.4) \quad \text{The maximal solution } r(t, \hat{t}, \hat{u}) \text{ of the scalar initial value problem } u' = g(t, u), u(\hat{t}) = \hat{u} \text{ exists and is bounded on } [\hat{t}, \infty) \text{ for each } (\hat{t}, \hat{u}) \in [t_0, \infty) \times \mathbb{R}^+.$$

Then, the largest interval of existence of any solution $x(t, \hat{t}, \hat{x})$ of (1)-(2) is $[\hat{t}, \infty)$ and $\|x(t, \hat{t}, \hat{x})\| \leq r(t, \hat{t}, \|\hat{x}\|)$ for all $t \in [\hat{t}, \infty)$.

Moreover, $\lim_{t \rightarrow \infty} x(t, \hat{t}, \hat{x})$ exists and is a finite vector in B for each

$$(\hat{t}, \hat{x}) \in [t_0, \infty) \times B.$$

It is also observed in [5] that if the equation $u' = -g(t, u)$, $u(\hat{t}) = \hat{u}$ had a maximal solution existing to the left and bounded on $[\ell, \hat{t}]$ then the interval of definition of $x(t, \hat{t}, \hat{x})$ can be extended to include $[\ell, \hat{t}]$ when (2.1), (2.2), and (2.3) of Theorem 2 hold.

From Theorem 2 we have conditions such that a solution of (1)-(2) satisfies the first half of asymptotic equilibrium. The following theorem begins our attack on the second half.

Theorem 3. If (1)-(2) satisfy the hypotheses of Theorem 2 then for $v \in B$ there exists a constant $T \in [t_0, \infty)$ and a sequence of functions $\{x_n(t)\}_{n=1}^{\infty}$ defined on $[T, \infty)$ and satisfying:

- (3.1) $\{x_n(t)\}_{n=1}^{\infty}$ is equicontinuous on $[T, \infty)$.
- (3.2) $\{x_n(t)\}_{n=1}^{\infty}$ is uniformly bounded on $[T, \infty)$, that is, there exists $M \in \mathbb{R}^+$ such that $\|x_n(t)\| \leq M$ for all $t \in [T, \infty)$ and for all n .
- (3.3) For each n we have $x_n(t)$ is a solution of $x' = f(t, x)$,
 $x(T + n) = v$.

Proof Let $r(t, t_0, \|v\|)$ be the maximal solution of $u' = g(t, u)$, $u(t_0) = \|v\|$ which by (2.4) exists and is bounded on $[t_0, \infty)$. Since $g(t, u) \geq 0$ we know $r(t)$ is non-decreasing so $\lim_{t \rightarrow \infty} r(t) = r(\infty) < \infty$ and

$$\|v\| \leq r(t) \leq r(\infty) \text{ for all } t \in [t_0, \infty).$$

For $\lambda \in \mathbb{R}^+$ let $u(t, t_0, \lambda)$ be the maximal solution of $u' = g(t, u)$, $u(t_0) = \lambda$. As before we have $\lim_{t \rightarrow \infty} u(t) = u(\infty) < \infty$ and $\lambda = u(t_0) \leq u(t) \leq u(\infty)$ for all $t \in [t_0, \infty)$. However by (2.3) $g(t, u)$ is non-decreasing

in u for each t so that

$$\infty > u(\infty) - u(t_0) = \int_{t_0}^{\infty} g(s, u(s)) \, ds \geq \int_{t_0}^{\infty} g(s, \lambda) \, ds.$$

Thus,

$$\int_{t_0}^{\infty} g(s, \lambda) \, ds < \infty \quad \text{for any } \lambda \in \mathbb{R}^+. \quad (3.4)$$

In particular, $\int_{t_0}^{\infty} g(s, 2r(\infty)) \, ds < \infty$ and so there exists $T \in [t_0, \infty)$

such that

$$\int_T^{\infty} g(s, 2r(\infty)) \, ds < r(\infty). \quad (3.5)$$

Applying Theorem 2 for each non-negative integer n we know there exists a solution, $x_n(t)$ of $x' = f(t, x)$, $x(T+n) = v$ on $[T+n, \infty)$ and

$$||x_n(t)|| < r_n(t, T+n, ||v||) \quad \text{on } [T+n, \infty) \quad (3.6)$$

where $r_n(t)$ is the maximal solution of $u' = g(t, u)$, $u(T+n) = ||v||$.

But $r_n(T+n) = ||v|| = r(t_0) \leq r(T+n)$, so by the Comparison Theorem [p.16,6] and combining with (3.6) we have

$$||x_n(t)|| \leq r(t) \leq r(\infty) \quad \text{on } [T+n, \infty). \quad (3.7)$$

Let $u_n(t, T+n, ||v||)$ be a solution to the left for $u' = -g(t, u)$, $u(T+n) = ||v||$. Now the largest interval of existence will contain $[T, T+n]$ unless u_n becomes unbounded. Suppose $u_n(t)$ becomes unbounded

and let $t_1, t_2 \in [T, T+n]$ with $t_1 < t_2$, $u_n(t_1) = 2r(\infty)$, $u_n(t_2) = r(\infty)$ and $2r(\infty) \geq u_n(t) \geq r(\infty)$ for all $t \in [t_1, t_2]$. Now

$$\begin{aligned} r(\infty) &= 2r(\infty) - r(\infty) = u_n(t_1) - u_n(t_2) \\ &= \left| \int_{t_1}^{t_2} g(s, u_n(s)) \, ds \right| = \int_{t_1}^{t_2} g(s, u(s)) \, ds \\ &\leq \int_{t_1}^{t_2} g(s, 2r(\infty)) \, ds \leq \int_T^\infty g(s, 2r(\infty)) \, ds < r(\infty). \end{aligned}$$

Hence, for each n we know $u_n(t)$ remains bounded by $2r(\infty)$ on $[T, T+n]$. Thus, for each n the maximal solution of $u' = -g(t, u)$, $u(T+n) = ||v||$ to the left exists and is bounded by $2r(\infty)$ on $[T, T+n]$ and by the remarks following Theorem 2 we know the interval of definition of $x_n(t)$ can be extended to $[T, \infty)$. Moreover, by combining (3.7) we have

$$||x_n(t)|| \leq 2r(\infty) \quad \text{for } t \in [T, \infty) \text{ and for each } n. \quad (3.8)$$

To complete the proof of this theorem we need to show $\{x_n(t)\}_{n=1}^\infty$ is equicontinuous. Let $\varepsilon > 0$ and note that by (3.5)

$$\int_T^\infty g(s, 2r(\infty)) \, ds < \infty \quad \text{and hence there exists } S \in [T, \infty) \text{ such that}$$

$\int_S^\infty g(s, 2r(\infty)) ds < \epsilon$. On the interval $[T, S+1] \times [0, 2r(\infty)]$ the continuous function $g(t, u)$ is bounded by a positive number L . Now if $|t_1 - t_2| < \min(1, \epsilon/L)$ then

$$\begin{aligned} ||x_n(t_1) - x_n(t_2)|| &= \left| \int_{t_1}^{t_2} f(s, x_n(s)) ds \right| \\ &\leq \left| \int_{t_1}^{t_2} g(s, ||x_n(s)||) ds \right| \leq \left| \int_{t_1}^{t_2} g(s, 2r(\infty)) ds \right|. \end{aligned}$$

However, if $t_1, t_2 \in [S, \infty)$ we have

$$\left| \int_{t_1}^{t_2} g(s, 2r(\infty)) ds \right| \leq \int_S^\infty g(s, 2r(\infty)) ds < \epsilon \text{ and if } t_1, t_2 \in [T, S+1]$$

$$\text{then } \left| \int_{t_1}^{t_2} g(s, 2r(\infty)) ds \right| \leq L|t_1 - t_2| < L \epsilon/L = \epsilon. \text{ Thus } \{x_n(t)\}_{n=1}^\infty$$

is equicontinuous.

We have most of the machinery needed to complete the proof of asymptotic equilibrium. If somehow we could conclude that for each $t \in [T, \infty)$ (or in fact any infinite subinterval of $[T, \infty)$ that $\{x_n(t)\}_{n=1}^\infty$

is relatively compact) we could apply Ascoli's Theorem [p. 155,8] and, with only a little work, be finished. For example in R^n since $\{x_n(t)\}_{n=1}^{\infty}$ is a bounded set then it is relatively compact. For arbitrary Banach spaces the authors in [5] assumed the additional hypothesis that f maps bounded sets into relatively compact sets. It is not clear that this condition successfully yields $\{x_n(t)\}_{n=1}^{\infty}$ being relatively compact for each $t \in [T, \infty)$. If we define $m(t) = \alpha(\{x_n(t) | n=1, 2, \dots\})$ we know from (1.4) of Lemma 1 that what we need is $m(t) = 0$ on $[T, \infty)$ or some infinite subinterval and we will give such conditions. To this end we consider the following.

Lemma 4. Suppose the hypotheses of Theorem 2 are satisfied. Let $v \in B$ and let $\{x_n(t)\}_{n=1}^{\infty}$ be the sequence of functions which exist by Theorem 3. For each $t \in [T, \infty)$ let $m(t) = \alpha(\{x_n(t)\}_{n=1}^{\infty})$. Then $m(t)$ is uniformly continuous on $[T, \infty)$.

Proof. Notice that $m(t_1) - m(t_2) = \alpha(\{x_n(t_1)\}_{n=1}^{\infty}) - \alpha(\{x_n(t_2)\}_{n=1}^{\infty}) \leq \alpha(\{x_n(t_1) - x_n(t_2)\}_{n=1}^{\infty})$ by (1.3). Using the equicontinuity of $\{x_n(t)\}_{n=1}^{\infty}$ and (1.5) the proof is complete.

Lemma 5. Suppose the hypotheses of Theorem 2 are satisfied and f is uniformly continuous on bounded subsets of $[T, \infty) \times B$. Let $\{x_n(t)\}_{n=1}^{\infty}$ be the sequence of functions which exist by Theorem 3. For $\hat{t} \in [T, \infty)$

and $h > 0$ we can express

$$x_n(\hat{t} + h) = x_n(\hat{t}) + hf(\hat{t}, x_n(\hat{t})) + h \epsilon_n(h) \quad \text{and}$$

$$x_n(\hat{t}) = x_n(\hat{t} - h) + hf(\hat{t}, x_n(\hat{t})) + h \tilde{\epsilon}_n(h) \quad \text{where}$$

$$\lim_{h \rightarrow 0^+} \alpha(\{\epsilon_n(h)\}_{n=1}^\infty) = 0 \quad \text{and} \quad \lim_{h \rightarrow 0^+} \alpha(\{\tilde{\epsilon}_n(h)\}_{n=1}^\infty) = 0.$$

Proof. We will give the proof of the first expression and note the other proof is similar. Let $\epsilon > 0$ and since f is uniformly continuous on $A = [\hat{t}, \hat{t} + 1] \times \{x \in B \mid \|x\| < 1\}$ there exists $\delta > 0$ such that $|t_1 - t_2| + \|y_1 - y_2\| < \delta$ implies $\|f(t_1, y_1) - f(t_2, y_2)\| < \epsilon$ for $(t_1, y_1), (t_2, y_2) \in A$. Since $\{x_n(t)\}_{n=1}^\infty$ is equicontinuous, there exists $\gamma > 0$ such that for $t \in (\hat{t} - \gamma, \hat{t} + \gamma)$ we have $\|x_n(t) - x_n(\hat{t})\| + |t - \hat{t}| < \delta$ for all n . Thus, for $h < \delta$ we know

$$\begin{aligned} \|h \epsilon_n(h)\| &= \|x_n(\hat{t} + h) - x_n(\hat{t}) - hf(\hat{t}, x_n(\hat{t}))\| \\ &= \left\| \int_{\hat{t}}^{\hat{t}+h} [f(s, x_n(s)) - f(\hat{t}, x_n(\hat{t}))] ds \right\| \leq \int_{\hat{t}}^{\hat{t}+h} \|f(s, x_n(s)) \\ &\quad - f(\hat{t}, x_n(\hat{t}))\| ds \leq \int_{\hat{t}}^{\hat{t}+h} \epsilon ds = \epsilon h. \end{aligned}$$

Thus, $\|\epsilon_n(h)\| \leq \epsilon$ for each n and by (1.5) we have $\alpha(\{\epsilon_n(h)\}_{n=1}^\infty) \leq 2\epsilon$

and consequently $\lim_{h \rightarrow 0^+} \alpha(\{\epsilon_n(h)\}_{n=1}^\infty) = 0$.

Theorem 6. Suppose the hypotheses of Theorem 2 and also that

(6.1) f is uniformly continuous on bounded subsets of $[T, \infty) \times B$.

(6.2) There exists $t^* \in [T, \infty)$ such that $m(t^*) = 0$.

(6.3) $G \in C[[T, \infty) \times \mathbb{R}^+, \mathbb{R}^+]$ such that $G(t, 0) = 0$ and the only solution $u(t)$ of $u' = G(t, u)$, $u(\hat{t}) = 0$ with $\lim_{t \rightarrow \hat{t}} \frac{u(t)}{t - \hat{t}} = 0$ is $u(t) \equiv 0$.

(6.4) For $h > 0$ and A a bounded subset of B we have

$$\alpha(\{x + hf(t, x) | x \in A\}) - \alpha(A) \leq hG(t, \alpha(A)).$$

Then $m(t) \equiv 0$ on $[t^*, \infty)$.

Proof. Recall $m(t) = \alpha(\{x_n(t)\}_{n=1}^\infty)$. Now using Lemma 5 and considering $h > 0$ we have

$$\begin{aligned} \frac{m(t+h) - m(t)}{h} &= \frac{\alpha(\{x_n(t+h)\}_{n=1}^\infty) - \alpha(\{x_n(t)\}_{n=1}^\infty)}{h} \\ &= \frac{\alpha(\{x_n(t) + hf(t, x_n(t)) - h\epsilon_n(h)\}_{n=1}^\infty) - \alpha(\{x_n(t)\}_{n=1}^\infty)}{h} \\ &\leq \frac{\alpha(\{x_n(t) + hf(t, x_n(t))\}_{n=1}^\infty) - \alpha(\{x_n(t)\}_{n=1}^\infty) + \alpha(\{h\epsilon_n(h)\}_{n=1}^\infty)}{h}. \end{aligned}$$

Now, using (6.4) and (1.2) we have

$$\begin{aligned} \frac{m(t+h) - m(t)}{h} &\leq \frac{hG\left(t, \alpha(\{x_n(t)\}_{n=1}^{\infty})\right) + h\alpha(\{\epsilon_n(h)\}_{n=1}^{\infty})}{h} \\ &= G\left(t, \alpha(\{x_n(t)\}_{n=1}^{\infty})\right) + \alpha(\{\epsilon_n(h)\}_{n=1}^{\infty}). \end{aligned}$$

Hence, by applying Lemma 5 we obtain $D_+m(t) \leq G(t, m(t))$. Again, by the Comparison Theorem [p. 19,6] since $D_+m(t) \leq G(t, m(t))$ on $[T, \infty)$ and $m(t^*) = 0$ we have $m(t)$ is less than or equal to the maximal solution of $u' = G(t, u)$, $u(t^*) = 0$.

Consider $\frac{x_n(t^* + h)}{h}$ and using Lemma 5 then for $h > 0$ we have

$$\begin{aligned} \frac{x_n(t^* + h)}{h} &= \frac{x_n(t^*) + hf(t^*, x_n(t^*)) + h\epsilon_n(h)}{h} \\ &= \frac{x_n(t^*) + hf(t^*, x_n(t^*))}{h} + \epsilon_n(h). \end{aligned}$$

Thus,

$$\begin{aligned} 0 &\leq \lim_{t \rightarrow t^*} \frac{m(t)}{t - t^*} = \lim_{t \rightarrow t^*} \frac{\alpha(\{x_n(t)\}_{n=1}^{\infty})}{t - t^*} = \lim_{h \rightarrow 0^+} \frac{\alpha(\{x_n(t^* + h)\}_{n=1}^{\infty})}{h} \\ &= \lim_{h \rightarrow 0^+} \frac{\alpha(\{x_n(t^*) + hf(t^*, x_n(t^*))\}_{n=1}^{\infty})}{h} + \lim_{h \rightarrow 0^+} \alpha(\{\epsilon_n(h)\}_{n=1}^{\infty}) \end{aligned}$$

$$\begin{aligned}
&\leq \lim_{h \rightarrow 0^+} \frac{hG(t^*, \alpha(\{x_n(t)\}_{n=1}^\infty))}{h} + 0 = G(t^*, \alpha(\{x_n(t^*)\}_{n=1}^\infty)) \\
&= G(t^*, m(t^*)) = G(t^*, 0) = 0.
\end{aligned}$$

Hence $\lim_{t \rightarrow t^*} \frac{m(t)}{t - t^*} = 0$ so by (6.3) we have $m(t) = 0$ for all $t \in [t^*, \infty)$.

Remark. It is proved in [7] that the system (1)-(2) has a local solution when f satisfies (6.1), (6.2), (6.3) and (6.4). Thus, we have sufficient hypotheses to assure local existence.

In Theorem 6 we assumed the existence of $t^* \in [T, \infty)$ such that $m(t^*) = 0$. Theorem 7 yields sufficient conditions to assure such a t^* exists.

Theorem 7. Suppose the hypotheses of Theorem 2 and also

(7.1) f is uniformly continuous on bounded subsets of $[T, \infty) \times B$.

(7.2) $F \in C([T, \infty) \times R^+, R^+]$ and $u' = F(t, u)$, $u(\hat{t}) = \hat{u}$ has a solution to the left for each $(\hat{t}, \hat{u}) \in [T, \infty) \times R^+$ and for any solution $u(t)$ there exists $t_1 \in (T, \hat{t}]$ such that $u(t_1) = 0$.

(7.3) For $h < 0$, A a bounded subset of B , and $t \in [T, \hat{t}]$ we have $\alpha(\{x + hf(t, x) \mid x \in A\}) - \alpha(A) \leq hF(t, \alpha(A))$.

Then, there exists a $t^* \in [T, \infty)$ such that $m(t^*) = 0$.

Proof. Recall $m(t) = \alpha(\{x_n(t)\}_{n=1}^{\infty})$ and from Lemma 4 we know $m(t)$ is continuous. Now using Lemma 5 and considering $h > 0$ we have

$$\begin{aligned} \frac{m(t) - m(t-h)}{h} &= \frac{\alpha(\{x_n(t)\}_{n=1}^{\infty}) - \alpha(\{x_n(t-h)\}_{n=1}^{\infty})}{h} \\ &= \frac{\alpha(\{x_n(t)\}_{n=1}^{\infty}) - \alpha(\{x_n(t) - hf(t, x_n(t)) - h\tilde{e}_n(h)\}_{n=1}^{\infty})}{h}. \end{aligned}$$

Now by (1.1) we have

$$\frac{m(t) - m(t-h)}{h} \geq \frac{\alpha(\{x_n(t)\}_{n=1}^{\infty}) - \alpha(\{x_n(t) - hf(t, x_n(t))\}_{n=1}^{\infty}) + \alpha(\{-h\tilde{e}_n(h)\}_{n=1}^{\infty})}{h}.$$

Next using (7.3) and (1.2) we have

$$\begin{aligned} \frac{m(t) - m(t-h)}{h} &\geq \frac{hF(t, \alpha(\{x_n(t)\}_{n=1}^{\infty})) + h\alpha(\{\tilde{e}_n(h)\}_{n=1}^{\infty})}{h} \\ &= F(t, \alpha(\{x_n(t)\}_{n=1}^{\infty})) + \alpha(\{\tilde{e}_n(h)\}_{n=1}^{\infty}). \end{aligned}$$

Thus,

$$D_-m(t) = \lim_{h \rightarrow 0^+} \frac{m(t) - m(t-h)}{h} \geq \lim_{h \rightarrow 0^+} F(t, \alpha(\{x_n(t)\}_{n=1}^{\infty})) + \lim_{h \rightarrow 0^+} \alpha(\{\tilde{e}_n(h)\}_{n=1}^{\infty}).$$

Since $\lim_{h \rightarrow 0^+} \alpha(\{x_n(h)\}_{n=1}^\infty) = 0$ by Lemma 5 then $D_m(t) = F(t, m(t))$.

From (3.8) we have $\|x_n(t)\| \leq 2r(\infty)$ on $[T, \infty)$ for any n so by (1.5) it follows that $m(t) \leq 4r(\infty)$. For $\hat{t} \in (T, \infty)$ let $R(t, \hat{t}, 4r(\infty))$ be the maximal solution to the left for $u' = F(t, u)$, $u(\hat{t}) = 4r(\infty)$. The existence of $R(t, \hat{t}, 4r(\infty))$ on $[T, \hat{t}]$ follows from (7.2). By the Comparison Theorem [p. 15, 6] since $D_m(t) \geq F(t, m(t))$ we have $m(t) \leq R(t, \hat{t}, 4r(\infty))$ on $[T, \hat{t}]$. However, by (7.1) there exists $t_1 \in (T, \hat{t}]$ with $R(t_1) = 0$ and so there exists $t^* \in [t_1, t]$ with $\alpha(\{x_n(t^*)\}_{n=1}^\infty) = m(t^*) = 0$ which means $\{x_n(t^*)\}_{n=1}^\infty$ is relatively compact.

Theorem 8. Suppose the hypotheses of Theorems 2, 6 and 7 are satisfied. Then (1)-(2) has asymptotic equilibrium.

Proof. The sequence $\{x_n(t)\}_{n=1}^\infty$ constructed in Theorem 3 has the property that $\{x_n(t)\}_{n=1}^\infty$ is relatively compact for each $t \in [t^*, \infty)$ by Theorems 6 and 7. Applying Ascoli's Theorem [p. 155, 8] we know there exists a subsequence which converges pointwise on $[t^*, \infty)$ and uniformly on compact subsets of $[t^*, \infty)$. To simplify the notation we will use $\{x_n(t)\}_{n=1}^\infty$ to denote the subsequence. Let $x(t) = \lim_{n \rightarrow \infty} x_n(t)$ for each $t \in [t^*, \infty)$ and note $x(t)$ is continuous by the uniform convergence on compact subsets. For each $t \in [t^*, \infty)$ we know from (3.3) that

$$x_n(t) = x_n(t^*) + \int_{t^*}^t f(s, x_n(s)) ds.$$

Since convergence is uniform on compact subsets then it follows that

$$x(t) = x(t^*) + \int_{t^*}^t f(s, x(s)) ds.$$

Thus, $x(t)$ is a solution of the system $x' = f(t, x)$, $x(t^*) = x(t^*)$.

It remains only to show that $\lim_{t \rightarrow \infty} x(t) = v$. Let $\epsilon > 0$ and then by (3.4) there exists an integer $S > t^*$ such that $\int_S^\infty g(s, 2r(\infty)) ds < \epsilon/2$.

If $\hat{t} \in [S, \infty)$ there exists N such that for all $n \geq N$ we know

$\|x_n(\hat{t}) - x(\hat{t})\| < \epsilon/2$. Consider $N + S$ and recall from (3.3) we have $x_{N+S}(T + N + S) = v$. Thus,

$$\begin{aligned} \|x(\hat{t}) - v\| &= \|x(\hat{t}) - x_{N+S}(T + N + S)\| \\ &\leq \|x(\hat{t}) - x_{N+S}(\hat{t})\| + \|x_{N+S}(\hat{t}) - x_{N+S}(T + N + S)\| \\ &\leq \epsilon/2 + \left\| \int_{\hat{t}}^{T+N+S} f(s, x_{N+S}(s)) ds \right\| \\ &\leq \epsilon/2 + \left| \int_{\hat{t}}^{T+N+S} g(s, \|x_{N+S}(s)\|) ds \right| \\ &\leq \epsilon/2 + \int_S^\infty g(s, 2r(\infty)) ds < \epsilon/2 + \epsilon/2 = \epsilon \end{aligned}$$

and so $\lim_{t \rightarrow \infty} x(t) = v$.

REFERENCES

1. Brauer, F., Global Behavior of Solutions of Ordinary Differential Equations, J. MATH. APPL., 2 (1961), pages 145-158.
2. Kuratowski, K., Topology, Vol. II, Academic Press, New York, 1962.
3. Ladas, G. and Lakshmikantham, V., Asymptotic Equilibrium of Ordinary Differential Systems, J. APPLICABLE ANAL. (to appear).
4. Ladas, G. and Lakshmikantham, V., Differential Equations in Abstract Spaces, Academic Press, New York, 1972.
5. Ladas, G. and Lakshmikantham, V., Global Existence and Asymptotic Equilibrium in Banach Spaces, J. INDIAN MATH. SOC., 36 (1972), pages 33-40.
6. Lakshmikantham, V. and Leela, S., Differential and Integral Inequalities Theory and Application, Vol. I., Academic Press, New York, 1969.
7. Li, Tien Yien, Existence of Solutions for Ordinary Differential Equations in Banach Spaces (to appear).
8. Royden, H. L., Real Analysis, MacMillan, New York, 1963.