DATE DUE

**CONSERVATIVE MOTION OF DISCRETE,
HEXAHEDRAL GYROSCOPE**

Donald Greenspan

TR # 315

CONSERVATIVE MOTION OF A DISCRETE, HEXAHEDRAL GYROSCOPE

Donald Greenspan

ABSTRACT:

Gyroscopic motion is simulated by applying a molecular dynamics formulation to a rigid hexahedron. The conservative dynamical differential equations are solved numerically in such a fashion that all the system invariants are preserved. Examples which included precession, nutation, and a combination of looping and cusp formation are described and discussed.

1. Introduction. Rigid body motion is of fundamental interest in mathematics, science, and engineering. In this paper we will study a simplistic approach to this area of study in the spirit of molecular mechanics. In particular, we will consider a discrete, hexahedral body whose six faces are congruent, regular triangles and simulate its motion as it spins like a gyroscope. The approach will not require the use of special coordinates, Cayley-Klein parameters, tensors, dyadics, or related concepts¹. All that will be required is three dimensional XYZ space and Newtonian mechanics. The numerical methodology will conserve exactly the same energy, linear momentum, and angular momentum as does the associated differential system. Computer examples, run in double precision on a 64 bit Alpha 275 personal computer, will be described and discussed.

2. A Discrete, Rigid Hexahedral Gyroscope. As shown in Figure 1, consider a hexahedron with six congruent, regular triangular faces. Let the edge length of each triangle be R and let the vertices be $P_i(x_i, y_i, z_i)$, $i = 1, 2, 3, 4, 5$. Initially, let

$$\begin{aligned}(x_1, y_1, z_1) &= (0, 0, 0), & (x_2, y_2, z_2) &= (0, R\sqrt{3}/3, R\sqrt{6}/3) \\(x_3, y_3, z_3) &= (\frac{1}{2}R, -R\sqrt{3}/6, R\sqrt{6}/3), & (x_4, y_4, z_4) &= (-\frac{1}{2}R, -R\sqrt{3}/6, R\sqrt{6}/3) \\(x_5, y_5, z_5) &= (0, 0, 2R\sqrt{6}/3)\end{aligned}$$

The geometric center $\bar{P} = (\bar{x}, \bar{y}, \bar{z})$ of the hexahedron is $(0, 0, R\sqrt{6}/3)$.

We define the neighbors of P_1 and P_5 to be P_2, P_3, P_4 . The neighbors of any one of P_2, P_3, P_4 are defined to be all the other vertices. thus, P_1 and P_5 have three neighbors each, while P_2, P_3 and P_4 have four.

Next, let us set P_2, P_3, P_4 in rotation in the plane $z = R\sqrt{6}/3$. As shown in Figure 2, let the velocity of each particle be perpendicular to the line joining that particle to the geometric center $\bar{P}$. Assume that P_2, P_3, P_4 each has the same speed V , so that we take the initial velocities $\vec{v}_i = (v_{ix}, v_{iy}, v_{iz})$, $i = 1, 2, 3, 4, 5$, of each P_i to be

$$\begin{aligned}\vec{v}_1 &= (0, 0, 0), \quad \vec{v}_2 = (V, 0, 0), \quad \vec{v}_3 = (-\frac{1}{2}V, -V\sqrt{3}/2, 0) \\ \vec{v}_4 &= (-\frac{1}{2}V, V\sqrt{3}/2, 0), \quad \vec{v}_5 = (0, 0, 0).\end{aligned}$$

Figure 1

Figure 2

Finally, we want the rotating hexahedron to be tilted, initially, relative to the Z axis, so that we assume that the line joining P_1 to the geometric center $\bar{P}$ forms an angle α relative to the Z axis. As shown in Figure 3, this will be done by rotating the XZ plane through an angle α . Thus, the new positions (x'_i, y'_i, z'_i) and the new velocities $(v_{ix'}, v_{iy'}, v_{iz'})$, $i = 1, 2, 3, 4, 5$, satisfy

$$\begin{aligned} x'_i &= x_i \cos \alpha + z_i \sin \alpha, \quad y'_i = y_i, \quad z'_i = -x_i \sin \alpha + z_i \cos \alpha, \\ v_{ix'} &= v_{ix} \cos \alpha + v_{iz} \sin \alpha, \quad v_{iy'} = v_{iy}, \quad v_{iz'} = -v_{ix} \sin \alpha + v_{iz} \cos \alpha. \end{aligned}$$

Thus, once the parameters R, V , and α are given, all initial data for a tilted, rotating, hexahedral gyroscope are determined.

3. Dynamical Equations. The motion of our rotating gyroscope is now treated as a five-body problem. At any time t , let P_i , $i = 1, 2, 3, 4, 5$, be located at $\vec{r}_i = (x_i, y_i, z_i)$, have velocity $\vec{v}_i = (\dot{x}_i, \dot{y}_i, \dot{z}_i) = (v_{ix}, v_{iy}, v_{iz})$, and have acceleration $\vec{a}_i = (\ddot{x}_i, \ddot{y}_i, \ddot{z}_i) = (\dot{v}_{ix}, \dot{v}_{iy}, \dot{v}_{iz})$. Let the mass of each P_i be m_i . For $i \neq j$, let $\vec{r}_{ij}$ be the vector from P_i to P_j and let r_{ij} be the magnitude of $\vec{r}_{ij}$, $i = 1, 2, 3, 4, 5$; $j = 1, 2, 3, 4, 5$; $i \neq j$. Let $\phi = \phi(r_{ij})$ be a potential function defined by the pair P_i, P_j , $i \neq j$. Then, for $i = 1, 2, 3, 4, 5$, the Newtonian dynamical equations for the motion of the particles are the second order differential equations.

$$(3.1) \quad m_i a_{ix} = -\frac{\partial \phi}{\partial r_{ij}} \frac{x_i - x_j}{r_{ij}} - \frac{\partial \phi}{\partial r_{ik}} \frac{x_i - x_k}{r_{ik}} - \frac{\partial \phi}{\partial r_{im}} \frac{x_i - x_m}{r_{im}} - \delta_i \frac{\partial \phi}{\partial r_{in}} \frac{x_i - x_n}{r_{in}}$$

$$(3.2) \quad m_i a_{iy} = -\frac{\partial \phi}{\partial r_{ij}} \frac{y_i - y_j}{r_{ij}} - \frac{\partial \phi}{\partial r_{ik}} \frac{y_i - y_k}{r_{ik}} - \frac{\partial \phi}{\partial r_{im}} \frac{y_i - y_m}{r_{im}} - \delta_i \frac{\partial \phi}{\partial r_{in}} \frac{y_i - y_n}{r_{in}}$$

$$(3.3) \quad m_i a_{iz} = -\frac{\partial \phi}{\partial r_{ij}} \frac{z_i - z_j}{r_{ij}} - \frac{\partial \phi}{\partial r_{ik}} \frac{z_i - z_k}{r_{ik}} - \frac{\partial \phi}{\partial r_{im}} \frac{z_i - z_m}{r_{im}} - \delta_i \frac{\partial \phi}{\partial r_{in}} \frac{z_i - z_n}{r_{in}} - g_i.$$

In (3.1)-(3.3), $\delta_1 = \delta_5 = 0$, $\delta_2 = \delta_3 = \delta_4 = 1$. Moreover, if $i = 1$, then $j = 2, k = 3, m = 4$; if $i = 2$, then $j = 1, k = 3, m = 4, n = 5$; if $i = 3$, then $j = 1, k = 2, m = 4, n = 5$; if $i = 4$, then $j = 1, k = 2, m = 3, n = 5$; if $i = 5$, then $j = 2, k = 3, m = 4$. Moreover, we choose $g_1 = 0$, $g_2 = g_3 = g_4 = g_5 > 0$, so that gravity acts only on P_2, P_3, P_4, P_5 .

Equations (3.1)-(3.3) are fully conservative, that is, they conserve system energy, linear momentum and angular momentum, which are called the system invariants¹.

4. Numerical Method. In order to solve system (3.1)-(3.3) in a fashion which conserves exactly the same energy, linear momentum and angular momentum, we first rewrite (3.1)-(3.3) as the following equivalent first order system:

$$(4.1) \quad \frac{dx_i}{dt} = v_{ix}$$

$$(4.2) \quad \frac{dy_i}{dt} = v_{iy}$$

$$(4.3) \quad \frac{dz_i}{dt} = v_{iz}$$

$$(4.4) \quad m_i \frac{dv_{ix}}{dt} = - \frac{\partial \phi}{\partial r_{ij}} \frac{x_i - x_j}{r_{ij}} - \frac{\partial \phi}{\partial r_{ik}} \frac{x_i - x_k}{r_{ik}} - \frac{\partial \phi}{\partial r_{im}} \frac{x_i - x_m}{r_{im}} - \delta_i \frac{\partial \phi}{\partial r_{in}} \frac{x_i - x_n}{r_{in}}$$

$$(4.5) \quad m_i \frac{dv_{iy}}{dt} = - \frac{\partial \phi}{\partial r_{ij}} \frac{y_i - y_j}{r_{ij}} - \frac{\partial \phi}{\partial r_{ik}} \frac{y_i - y_k}{r_{ik}} - \frac{\partial \phi}{\partial r_{im}} \frac{y_i - y_m}{r_{im}} - \delta_i \frac{\partial \phi}{\partial r_{in}} \frac{y_i - y_n}{r_{in}}$$

$$(4.6) \quad m_i \frac{dv_{iz}}{dt} = - \frac{\partial \phi}{\partial r_{ij}} \frac{z_i - z_j}{r_{ij}} - \frac{\partial \phi}{\partial r_{ik}} \frac{z_i - z_k}{r_{ik}} - \frac{\partial \phi}{\partial r_{im}} \frac{z_i - z_m}{r_{im}} - \delta_i \frac{\partial \phi}{\partial r_{in}} \frac{z_i - z_n}{r_{in}} - g_i.$$

We now choose difference equation approximations for (4.1)-(4.6). For a fixed time step Δt , let $t_N = N \Delta t$, $N = 0, 1, 2, 3, \dots$. At t_N let P_i be at $\vec{r}_{i,N} = (x_{i,N}, y_{i,N}, z_{i,N})$ with velocity $\vec{v}_{i,N} = (v_{i,x,N}, v_{i,y,N}, v_{i,z,N})$. Then (4.1)-(4.6) will be approximated by

$$(4.7) \quad \frac{x_{i,N+1} - x_{i,N}}{\Delta t} = \frac{v_{i,x,N+1} + v_{i,x,N}}{2}$$

$$(4.8) \quad \frac{y_{i,N+1} - y_{i,N}}{\Delta t} = \frac{v_{i,y,N+1} + v_{i,y,N}}{2}$$

$$(4.9) \quad \frac{z_{i,N+1} - z_{i,N}}{\Delta t} = \frac{v_{i,z,N+1} + v_{i,z,N}}{2}$$

$$(4.10) \quad m_i \frac{v_{i,x,N+1} - v_{i,x,N}}{\Delta t} = - \frac{\phi(r_{ij,N+1}) - \phi(r_{ij,N})}{r_{ij,N+1} - r_{ij,N}} \cdot \frac{x_{i,N+1} + x_{i,N} - x_{j,N+1} - x_{j,N}}{r_{ij,N+1} + r_{ij,N}} \\ - \frac{\phi(r_{ik,N+1}) - \phi(r_{ik,N})}{r_{ik,N+1} - r_{ik,N}} \cdot \frac{x_{i,N+1} + x_{i,N} - x_{k,N+1} - x_{k,N}}{r_{ik,N+1} + r_{ik,N}} \\ - \frac{\phi(r_{im,N+1}) - \phi(r_{im,N})}{r_{im,N+1} - r_{im,N}} \cdot \frac{x_{i,N+1} + x_{i,N} - x_{m,N+1} - x_{m,N}}{r_{im,N+1} + r_{im,N}} \\ - \delta_i \frac{\phi(r_{in,N+1}) - \phi(r_{in,N})}{r_{in,N+1} - r_{in,N}} \cdot \frac{x_{i,N+1} + x_{i,N} - x_{n,N+1} - x_{n,N}}{r_{in,N+1} + r_{in,N}}$$

$$(4.11) \quad m_i \frac{v_{i,y,N+1} - v_{i,y,N}}{\Delta t} = - \frac{\phi(r_{ij,N+1}) - \phi(r_{ij,N})}{r_{ij,N+1} - r_{ij,N}} \cdot \frac{y_{i,N+1} + y_{i,N} - y_{j,N+1} - y_{j,N}}{r_{ij,N+1} + r_{ij,N}} \\ - \frac{\phi(r_{ik,N+1}) - \phi(r_{ik,N})}{r_{ik,N+1} - r_{ik,N}} \cdot \frac{y_{i,N+1} + y_{i,N} - y_{k,N+1} - y_{k,N}}{r_{ik,N+1} + r_{ik,N}}$$

$$\begin{aligned}
& - \frac{\phi(r_{im,N+1}) - \phi(r_{im,N})}{r_{im,N+1} - r_{im,N}} \cdot \frac{y_{i,N+1} + y_{i,N} - y_{m,N+1} - y_{m,N}}{r_{im,N+1} + r_{im,N}} \\
& - \delta_i \frac{\phi(r_{in,N+1}) - \phi(r_{in,N})}{r_{in,N+1} - r_{in,N}} \cdot \frac{y_{i,N+1} + y_{i,N} - y_{n,N+1} - y_{n,N}}{r_{in,N+1} + r_{in,N}} \\
(4.12) \quad m_i \frac{v_{i,z,N+1} - v_{i,z,N}}{\Delta t} = & - \frac{\phi(r_{ij,N+1}) - \phi(r_{ij,N})}{r_{ij,N+1} - r_{ij,N}} \cdot \frac{z_{i,N+1} + z_{i,N} - z_{j,N+1} - z_{j,N}}{r_{ij,N+1} + r_{ij,N}} \\
& - \frac{\phi(r_{ik,N+1}) - \phi(r_{ik,N})}{r_{ik,N+1} - r_{ik,N}} \cdot \frac{z_{i,N+1} + z_{i,N} - z_{k,N+1} - z_{k,N}}{r_{ik,N+1} + r_{ik,N}} \\
& - \frac{\phi(r_{im,N+1}) - \phi(r_{im,N})}{r_{im,N+1} - r_{im,N}} \cdot \frac{z_{i,N+1} + z_{i,N} - z_{m,N+1} - z_{m,N}}{r_{im,N+1} + r_{im,N}} \\
& - \delta_i \frac{\phi(r_{in,N+1}) - \phi(r_{in,N})}{r_{in,N+1} - r_{in,N}} \cdot \frac{z_{i,N+1} + z_{i,N} - z_{n,N+1} - z_{n,N}}{r_{in,N+1} + r_{in,N}} \\
& - g_i
\end{aligned}$$

Difference equations (4.7)-(4.12) are consistent with differential equations (4.1)-(4.6) and conserve exactly the same system invariants^{2,3} as do (3.1)-(3.3). System (4.7)-(4.12), for each of $N = 0, 1, 2, 3, \dots$, consists of 30 equations in the unknowns $x_{i,N+1}$, $y_{1,N+1}$, $z_{i,N+1}$, $v_{i,x,N+1}$, $v_{i,y,N+1}$, $v_{i,z,N+1}$, $i = 1, 2, 3, 4, 5$; and in the knowns $x_{i,N}$, $y_{i,N}$, $z_{i,N}$, $v_{i,x,N}$, $v_{i,y,N}$, $v_{i,z,N}$. These are solved readily by Newton's method².

In all the examples to be described in the next section, the results will be given relative to P_1 , whose position and velocity coordinates at the end of each time step are set to $\vec{0}$. By this means, the motion of the hexahedron's geometric center $\vec{P}$ fully characterizes the motion of the gyroscope.

Typical FORTRAN programs for generating initial data and for trajectory calculation from given initial data are available for the interested reader in the Appendices of Greenspan⁴.

5. **Examples.** In considering examples, we must first choose a potential function ϕ . We do this in cgs units and in such a fashion that we ensure that the tetrahedron is rigid. To accomplish this, we introduce the following classical, molecular type function^{5,6}:

$$(5.1) \quad \phi = A \left[-\frac{1}{r_{ij}^3} + \frac{1}{r_{ij}^5} \right], \quad A > 0,$$

in which A is sufficiently large to impose rigidity. The choice of exponents in (5.1) prevents the explosive behavior characteristic of real molecules, yet allows for attractive and repulsive interaction, as is desirable for aggregates of molecules⁶.

From (5.1), it follows that the magnitude F of the force $\vec{F}$ determined by ϕ satisfies

$$F = A \left[-\frac{3}{r_{ij}^4} + \frac{5}{r_{ij}^6} \right].$$

Thus, $F(\bar{r}) = 0$ provided $\bar{r} = (5/3)^{1/2} \sim 1.290994449$.

We now choose the parameter R to be

$$(5.2) \quad R = 1.290994449..$$

For this value of R the force between any two neighboring particles is zero, so that the gyroscope is physically stable. The parameters A , g , and m_i are scaled for computational convenience to be $A = 10^6$, $g = 0.980$, $m_i = 1$, $i = 1, 2, 3, 4, 5$, unless otherwise indicated. The time step Δt is chosen to be $\Delta t = 10^{-5}$. In the figures to be given, units on the Z axis are often rescaled to accentuate the character of the resulting trajectory. In all the calculations, the distance between any two neighbors is always 1.291.

In this first example, let $V = 20$, $\alpha = 8^\circ$. Figure 4 shows the resulting circular precession of the geometric center every 10000 time steps through 28000000 time steps. The graph consists of 2800 points. $\vec{P}$ has completed just over one complete rotation. the $\bar{z}$ values vary in the narrow range $1.0435 < \bar{z} < 1.04395$. Figure 5 shows the projection in the XY plane of the motion shown in Figure 4.

In the next six examples, $V = 15$, but α varies successively through the values 15° , 30° , 45° , 60° , 75° , 90° . The results are shown in Figures 6-11. The figures reveal precession of the geometric center with the radius of the trajectory increasing with α . For $\alpha = 90^\circ$, one finds $-0.006 < \bar{z} < 0.000$.

For the next example, set $V = 3$, $\alpha = 16^\circ$. The resulting motion of the geometric center is the nutation shown in Figure 12. In this example $\bar{z}$ varies in the range $0.4 < \bar{z} < 0.54$, so that the gyroscope displays a periodic rising and falling motion. The projection of the trajectory in the XY plane is shown in Figure 13. Figures 12 and 13 indicate clearly the formation of cusps.

For our final example, choose again $V = 15$ and $\alpha = 15^\circ$, but let the mass m_2 of P_2 be 3. Thus, the resulting hexahedron, though symmetric, is nonhomogeneous. The resulting trajectory of $\vec{P}$ is shown every 10000 time steps through 30000000 time steps in Figure 14. $\vec{P}$ has completed just over one complete rotation in this time and the graph contains 3000 points. That this graph is different from all previous graphs is quite visible. To analyze the resulting behavior in greater detail, we first examined the projection of the trajectory in the XY plane, since the variation in $\bar{z}$ was entirely in the small range $1.0155 < \bar{z} < 1.0205$. This projection is shown in Figure 15 and reveals that the variations of X and Y must be relatively complex. To study these variations more precisely, the motion was simulated over its first 500000 steps and plotted every 500 time steps. The projection of the resulting 1000 points in the XY plane is shown in Figure 16. What appears to be happening is that the X and Y coordinates are looping. However, plotting every 500 points has resulted in a graph with polygonal shapes rather than smooth shapes. To analyze the motion further, we concentrated only on the lower right corner of

the graph show in Figure 16. For this purpose, we reran the trajectory but for only 150000 time steps and plotted every 50 time steps. The projected motion in the XY plane is shown in Figure 17 and contains 3000 points. What is revealed is both looping and cusp formation, which to us was totally unexpected.

6. Remarks. It would be most interesting to plot the trajectories shown in figures 14 and 15 with the accuracy used in Figure 17. However, to plot every 50 time steps through 30000000 time steps would require the ability to plot the resulting 600000 points. Our Delta Graph package did not have this capability. Further, one must be able to plot on a relatively large set of axes to exhibit the details of Figure 17 throughout the entire trajectory.

Though small perturbations of the parameters used for the results shown in Figures 4-13 resulted in entirely analogous results, the introduction of nonhomogeneity results in complexities which were unexpected. Though the motion of homogeneous, continuous gyroscopes can at times be analyzed mathematically, this does not seem to be the case when nonhomogeneity is introduced^{7,8}. Further study of examples like the final one in Section 5 are in progress, but the graphics difficulties and the large number of parameter choices under study are resulting in very slow progress.

Finally, note that any other classical molecular type function which ensures rigidity can be used in place of (5.1).

Figure 5. $V=20$, $\alpha=8^\circ$

Figure 4. $V=20$, $\alpha=8^\circ$

z

Figure 6. $V=15$, $\alpha=15^\circ$

z

Figure 7. $V=15$, $\alpha=30^\circ$

Figure 9. $V=15$, $\alpha=60^\circ$

Figure 8. $V=15$, $\alpha=45^\circ$

Figure 11, $V=15, \alpha$

Figure 10, $V=15, \alpha$

Figure 13. $V=3$, $\alpha=16^\circ$

Figure 12. $V=3$, $\alpha=16^\circ$

Figure 14. $V=15$, $\alpha=15^\circ$, $m_2=3$

Figure 15. $V=15$, $\alpha=15^\circ$, $m_2=3$

Figure 17. $V=15$, $\alpha=15^\circ$, $m_2=3$.

Figure 16. $V=15$, $\alpha=15^\circ$, $m_2=3$.

REFERENCES

1. H. Goldstine, CLASSICAL MECHANICS, 2nd Edition, Addison-Wesley, Reading, 1980, Chaps. iv, v.
2. D. Greenspan, ARITHMETIC APPLIED MATHEMATICS, Pergamon, Oxford, 1980.
3. D. Greenspan, "Completely Conservative, Covariant Numerical Methodology", Comp. Math. Applic., 29, 1995, p. 37.
4. D. Greenspan, TR 315, Math. Dept., UT Arlington, Arlington, Texas, 1996.
5. R. P. Feynman, R. B. Leighton, and M. Sands, THE FEYNMAN LECTURES ON PHYSICS, vol. I, Addison-Wesley, Reading, 1963.
6. D. Greenspan, QUASIMOLECULAR MODELLING, World Sci. Press, Singapore, 1991.
7. L. D. Landau and E. M. Lifshitz, MECHANICS (3rd Edition), Pergamon, Oxford, 1976.
8. A. Gray, GYROSTATICS AND ROTATIONAL MOTION, Dover, N.Y., 1959.

C THE PRIME COORDINATES.

```

X1P=X1*COS(ALPHA)+Z1*SIN(ALPHA)
Y1P=Y1
Z1P=-X1*SIN(ALPHA)+Z1*COS(ALPHA)
X2P=X2*COS(ALPHA)+Z2*SIN(ALPHA)
Y2P=Y2
Z2P=-X2*SIN(ALPHA)+Z2*COS(ALPHA)
X3P=X3*COS(ALPHA)+Z3*SIN(ALPHA)
Y3P=Y3
Z3P=-X3*SIN(ALPHA)+Z3*COS(ALPHA)
X4P=X4*COS(ALPHA)+Z4*SIN(ALPHA)
Y4P=Y4
Z4P=-X4*SIN(ALPHA)+Z4*COS(ALPHA)
X5P=X5*COS(ALPHA)+Z5*SIN(ALPHA)
Y5P=Y5
Z5P=-X5*SIN(ALPHA)+Z5*COS(ALPHA)
VX1P=VX1*COS(ALPHA)+VZ1*SIN(ALPHA)
VY1P=VY1
VZ1P=-VX1*SIN(ALPHA)+VZ1*COS(ALPHA)
VX2P=VX2*COS(ALPHA)+VZ2*SIN(ALPHA)
VY2P=VY2
VZ2P=-VX2*SIN(ALPHA)+VZ2*COS(ALPHA)
VX3P=VX3*COS(ALPHA)+VZ3*SIN(ALPHA)
VY3P=VY3
VZ3P=-VX3*SIN(ALPHA)+VZ3*COS(ALPHA)
VX4P=VX4*COS(ALPHA)+VZ4*SIN(ALPHA)
VY4P=VY4
VZ4P=-VX4*SIN(ALPHA)+VZ4*COS(ALPHA)
VX5P=VX5*COS(ALPHA)+VZ5*SIN(ALPHA)
VY5P=VY5
VZ5P=-VX5*SIN(ALPHA)+VZ5*COS(ALPHA)
X1=X1P
Y1=Y1P
Z1=Z1P
X2=X2P
Y2=Y2P
Z2=Z2P
X3=X3P
Y3=Y3P
Z3=Z3P
X4=X4P
Y4=Y4P
Z4=Z4P
X5=X5P
Y5=Y5P
Z5=Z5P
XBARNEW=(X1+X2+X3+X4+X5)/5.
YBARNEW=(Y1+Y2+Y3+Y4+Y5)/5.
ZBARNEW=(Z1+Z2+Z3+Z4+Z5)/5.
VX1=VX1P
VY1=VY1P
VZ1=VZ1P
VX2=VX2P
VY2=VY2P
VZ2=VZ2P
VX3=VX3P
VY3=VY3P
VZ3=VZ3P
VX4=VX4P
VY4=VY4P

```

VZ4=VZ4P

VX5=VX5P

VY5=VY5P

VZ5=VZ5P

WRITE (22,1001) X1,Y1,Z1,VX1,VY1,VZ1

WRITE (22,1001) X2,Y2,Z2,VX2,VY2,VZ2

WRITE (22,1001) X3,Y3,Z3,VX3,VY3,VZ3

WRITE (22,1001) X4,Y4,Z4,VX4,VY4,VZ4

WRITE (22,1001) X5,Y5,Z5,VX5,VY5,VZ5

WRITE (22,9999) YBAR,VBAR,ZBAR

WRITE (22,9999) YBARNEW,YBARNEW,ZBARNEW

2000 FORMAT (3F15,10)

1001 FORMAT (3F15,10,3F15,10)

23 STOP

END

C IT IS IMPORTANT TO NOTE THAT THE IMPLICIT METHOD TO
C BE USED CONSERVES ENERGY, LINEAR MOMENTUM, AND ANGULAR
C MOMENTUM.

DOUBLE PRECISION XO(5),YO(5),VO(5),VYO(5),

IX(5),Y(5),VX(5),VY(5),

IXX(5),YY(5),VXX(5),VYY(5)

IZO(5),VZO(5),Z(5),VZ(5),ZZ(5),VZZ(5)

IR(5),RN(5),RR(5),XXX(5),YYY(5),ZZZ(5),S)

IXBAR,YBAR,ZBAR

IXBAR,YBAR,ZBAR

OPEN (UNIT=21,FILE='HEXAL.DAT',STATUS='OLD')

OPEN (UNIT=22,FILE='HEXAL.OUT',STATUS='NEW')

OPEN (UNIT=23,FILE='HEXAL.XKE',STATUS='NEW')

OPEN (UNIT=24,FILE='HEXAL.BAR',STATUS='NEW')

AF=0.00000

WPRINT=1.0000

K=1

DELT=0.00001

TIME=0.0

C READ IN THE INITIAL DATA.

READ (21,10) (XO(I),YO(I),VO(I),VYO(I),VZO(I),

VI=1.5)

FORMAT(3F15.10,3F15.10)

C WE NOW SOLVE FOR THE NEW POSITIONS AND VELOCITIES WITH

C FORMULAS WHICH REQUIRE ITERATION. WE WILL USE THE

C IDEAS OF THE GENERALIZED NEWTON'S METHOD WITHOUT THE

C UPDATING EACH ITERATE.

C SET THE INPUT DATA.

DO 30 I=1,5

X(I,1)=XO(I)

Y(I,1)=YO(I)

Z(I,1)=ZO(I)

VX(I,1)=VXO(I)

VY(I,1)=VYO(I)

VZ(I,1)=VZO(I)

CONTINUE

DO 10 I=1,5

X(I,1)=X(I,2)

Y(I,1)=Y(I,2)

Z(I,1)=Z(I,2)

VX(I,1)=VX(I,2)

VY(I,1)=VY(I,2)

VZ(I,1)=VZ(I,2)

CONTINUE

C FIX THE FIRST GUESS X(I,2) OF THE ITERATION.

DO 1633 I=1,5

X(I,2)=X(I,1)

Y(I,2)=Y(I,1)

Z(I,2)=Z(I,1)

VX(I,2)=VX(I,1)

VY(I,2)=VY(I,1)

VZ(I,2)=VZ(I,1)

CONTINUE

DO 103 I=1,5

DO 102 J=1,5

R(I,J)=(X(I,1)-X(J,1))*Z+(Y(I,1)-Y(J,1))*Z

R(I,J)=OSQRT(R(I,J))

CONTINUE

102

```

103     CONTINUE
      XTIME=XTIME+DELT
      KK=1
      GO TO 110
105     DO 108 JJ=1,5
      X(JJ,2)=X(JJ,3)
      Y(JJ,2)=Y(JJ,3)
      Z(JJ,2)=Z(JJ,3)
      VX(JJ,2)=VX(JJ,3)
      P    VY(JJ,2)=VY(JJ,3)
      VZ(JJ,2)=VZ(JJ,3)
108     CONTINUE
110     DO 203 I=1,5
      DO 202 J=1,5
      RN(I,J)=(X(I,2)-X(J,2))**2+(Y(I,2)-Y(J,2))**2
      + (Z(I,2)-Z(J,2))**2
      RN(I,J)=DSQRT(RN(I,J))
202     CONTINUE
203     CONTINUE
      DO 109 I=1,5
      X(I,3)=X(I,1)+0.5*DELT*(VX(I,2)+VX(I,1))
      Y(I,3)=Y(I,1)+0.5*DELT*(VY(I,2)+VY(I,1))
      Z(I,3)=Z(I,1)+0.5*DELT*(VZ(I,2)+VZ(I,1))
109     CONTINUE
      DO 300 I=1,5
      DO 299 J=1,5
      XXX(I,J)=X(I,3)+X(I,1)-X(J,3)-X(J,1)
      YYY(I,J)=Y(I,3)+Y(I,1)-Y(J,3)-Y(J,1)
      ZZZ(I,J)=Z(I,3)+Z(I,1)-Z(J,3)-Z(J,1)
299     CONTINUE
300     CONTINUE
      DO 400 I=1,4
      DO 399 J=I+1,5
      RR(I,J)=-A*(R(I,J)**4+(R(I,J)**3)*RN(I,J)
      + (R(I,J)**2)*(RN(I,J)**2)+R(I,J)*(RN(I,J)**3)
      + (RN(I,J)**4)-(R(I,J)**2)*(RN(I,J)**2)*
      + (RN(I,J)**2+RN(I,J)*R(I,J)+R(I,J)**2))/
      + ((RN(I,J)**5)*(R(I,J)**5)*(RN(I,J)+R(I,J)))
399     CONTINUE
400     CONTINUE
      VX(1,3)=VX(1,1)-DELT*(XXX(1,2)*RR(1,2) +XXX(1,3)*RR(1,3)
      +XXX(1,4)*RR(1,4))
      VY(1,3)=VY(1,1)-DELT*(YYY(1,2)*RR(1,2) +YYY(1,3)*RR(1,3)
      +YYY(1,4)*RR(1,4))
      VZ(1,3)=VZ(1,1)-DELT*(ZZZ(1,2)*RR(1,2) +ZZZ(1,3)*RR(1,3)
      +ZZZ(1,4)*RR(1,4))
C NOTE THAT P1 IS INDEPENDENT OF GRAVITY.
      VX(2,3)=VX(2,1)-DELT*(XXX(2,1)*RR(1,2) +XXX(2,3)*RR(2,3)
      +XXX(2,4)*RR(2,4)+XXX(2,5)*RR(2,5))
      VY(2,3)=VY(2,1)-DELT*(YYY(2,1)*RR(1,2) +YYY(2,3)*RR(2,3)
      +YYY(2,4)*RR(2,4)+YYY(2,5)*RR(2,5))
      VZ(2,3)=VZ(2,1)-DELT*(ZZZ(2,1)*RR(1,2) +ZZZ(2,3)*RR(2,3)
      +ZZZ(2,4)*RR(2,4)+ZZZ(2,5)*RR(2,5))
      1-DELT*.98
      VX(3,3)=VX(3,1)-DELT*(XXX(3,1)*RR(1,3) +XXX(3,2)*RR(2,3)
      +XXX(3,4)*RR(3,4)+XXX(3,5)*RR(3,5))
      VY(3,3)=VY(3,1)-DELT*(YYY(3,1)*RR(1,3) +YYY(3,2)*RR(2,3)
      +YYY(3,4)*RR(3,4)+YYY(3,5)*RR(3,5))
      VZ(3,3)=VZ(3,1)-DELT*(ZZZ(3,1)*RR(1,3) +ZZZ(3,2)*RR(2,3)
      +ZZZ(3,4)*RR(3,4)+ZZZ(3,5)*RR(3,5))

```

```

1-DELT=.98
VX(4,3)=VX(4,1)-DELT*(XXX(4,1)*RR(1,4)+XXX(4,2)*RR(2,4)
+XXX(4,3)*RR(3,4)+XXX(4,5)*RR(4,5))
VY(4,3)=VY(4,1)-DELT*(YYY(4,1)*RR(1,4)+YYY(4,2)*RR(2,4)
+YYY(4,3)*RR(3,4)+YYY(4,5)*RR(4,5))
VZ(4,3)=VZ(4,1)-DELT*(ZZZ(4,1)*RR(1,4)+ZZZ(4,2)*RR(2,4)
+ZZZ(4,3)*RR(3,4)+ZZZ(4,5)*RR(4,5))
1-DELT=.98
VX(5,3)=VX(5,1)-DELT*(XXX(5,2)*RR(2,5)
+XXX(5,3)*RR(3,5)+XXX(5,4)*RR(4,5))
VY(5,3)=VY(5,1)-DELT*(YYY(5,2)*RR(2,5)
+YYY(5,3)*RR(3,5)+YYY(5,4)*RR(4,5))
VZ(5,3)=VZ(5,1)-DELT*(ZZZ(5,2)*RR(2,5)
+ZZZ(5,3)*RR(3,5)+ZZZ(5,4)*RR(4,5))
1-DELT=.98
KK=KK+1
IF (KK.GT.50000) GO TO 33
DO 9875 I=1,5
  XX(I)=ABS(X(I,3)-X(I,2))
  YY(I)=ABS(Y(I,3)-Y(I,2))
  ZZ(I)=ABS(Z(I,3)-Z(I,2))
  VXX(I)=ABS(VX(I,3)-VX(I,2))
  VYY(I)=ABS(VY(I,3)-VY(I,2))
  VZZ(I)=ABS(VZ(I,3)-VZ(I,2))
9875  CONTINUE
  INDEX=-1
  DO 500 I=1,5
    IF (XX(I).GT.0.000000000005) INDEX=+1
    IF (YY(I).GT.0.000000000005) INDEX=+1
    IF (ZZ(I).GT.0.000000000005) INDEX=+1
    IF (VXX(I).GT.0.000000000005) INDEX=+1
    IF (VYY(I).GT.0.000000000005) INDEX=+1
    IF (VZZ(I).GT.0.000000000005) INDEX=+1
500  CONTINUE
    IF (INDEX.EQ.+1) GO TO 105
    DO 1009 JJ=1,5
      X(JJ,2)=X(JJ,3)
      Y(JJ,2)=Y(JJ,3)
      Z(JJ,2)=Z(JJ,3)
      VX(JJ,2)=VX(JJ,3)
      VY(JJ,2)=VY(JJ,3)
      VZ(JJ,2)=VZ(JJ,3)
1009  CONTINUE
C WE NOW DETERMINE DATA RELATIVE TO P1.
X(2,2)=X(2,2)-X(1,2)
Y(2,2)=Y(2,2)-Y(1,2)
Z(2,2)=Z(2,2)-Z(1,2)
X(3,2)=X(3,2)-X(1,2)
Y(3,2)=Y(3,2)-Y(1,2)
Z(3,2)=Z(3,2)-Z(1,2)
X(4,2)=X(4,2)-X(1,2)
Y(4,2)=Y(4,2)-Y(1,2)
Z(4,2)=Z(4,2)-Z(1,2)
X(5,2)=X(5,2)-X(1,2)
Y(5,2)=Y(5,2)-Y(1,2)
Z(5,2)=Z(5,2)-Z(1,2)
X(1,2)=0.0
Y(1,2)=0.0
Z(1,2)=0.0
VX(2,2)=VX(2,2)-VX(1,2)

```

```

VY(2,2)=VY(2,2)-VY(1,2)
VZ(2,2)=VZ(2,2)-VZ(1,2)
VX(3,2)=VX(3,2)-VX(1,2)
VY(3,2)=VY(3,2)-VY(1,2)
VZ(3,2)=VZ(3,2)-VZ(1,2)
VX(4,2)=VX(4,2)-VX(1,2)
VY(4,2)=VY(4,2)-VY(1,2)
VZ(4,2)=VZ(4,2)-VZ(1,2)
VX(5,2)=VX(5,2)-VX(1,2)
VY(5,2)=VY(5,2)-VY(1,2)
VZ(5,2)=VZ(5,2)-VZ(1,2)
VX(1,2)=0.0
VY(1,2)=0.0
VZ(1,2)=0.

```

```

K=K+1

```

```

IF (MOD(K,KPRINT),GT.0) GO TO 82

```

```

V1SQ=VX(1,2)*VX(1,2)+VY(1,2)*VY(1,2)+VZ(1,2)*VZ(1,2)
V2SQ=VX(2,2)*VX(2,2)+VY(2,2)*VY(2,2)+VZ(2,2)*VZ(2,2)
V3SQ=VX(3,2)*VX(3,2)+VY(3,2)*VY(3,2)+VZ(3,2)*VZ(3,2)
V4SQ=VX(4,2)*VX(4,2)+VY(4,2)*VY(4,2)+VZ(4,2)*VZ(4,2)
V5SQ=VX(5,2)*VX(5,2)+VY(5,2)*VY(5,2)+VZ(5,2)*VZ(5,2)
ENERGY=-A/((R(1,2)**3))-A/((R(1,3)**3))
!-A/((R(1,4)**3))-A/((R(2,3)**3))
!-A/((R(2,4)**3))-A/((R(2,5)**3))
!-A/((R(3,4)**3))-A/((R(3,5)**3))
!-A/((R(4,5)**3))
!+A/((R(1,2)**5))+A/((R(1,3)**5))
!+A/((R(1,4)**5))+A/((R(2,3)**5))
!+A/((R(2,4)**5))+A/((R(2,5)**5))+A/((R(3,4)**5))
!+A/((R(3,5)**5))+A/((R(4,5)**5))
!+.98*(Z(2,2)+Z(3,2)+Z(4,2)+Z(5,2))

```

```

ENERGY=ENERGY+(V1SQ+V2SQ+V3SQ+V4SQ+V5SQ)

```

```

ENERGY=.000001*ENERGY

```

```

WRITE (23,8000) K,ENERGY,RN(1,2),RN(1,3),RN(1,4),

```

```

RN(2,3),RN(2,4),RN(3,4),RN(2,5),RN(3,5),RN(4,5),XTIME

```

```

8000 FORMAT (I13,F12.6,9F7.3,F12.5)

```

```

XBAR=(X(1,2)+X(2,2)+X(3,2)+X(4,2)+X(5,2))/5.

```

```

YBAR=(Y(1,2)+Y(2,2)+Y(3,2)+Y(4,2)+Y(5,2))/5.

```

```

ZBAR=(Z(1,2)+Z(2,2)+Z(3,2)+Z(4,2)+Z(5,2))/5.

```

C NOTE THAT XBAR,YBAR,ZBAR ARE RELATIVE TO P1 SO THAT

C WE CAN EASILY PLOT THE MOTION OF THE MASS CENTER.

```

WRITE (24,938) XBAR,YBAR,ZBAR

```

```

938 FORMAT (3F16.10)

```

```

82 IF (K.LE.10000000) GO TO 65

```

```

9383 WRITE (22,10) (X(I,2),Y(I,2),Z(I,2),VX(I,2),

```

```

VY(I,2),VZ(I,2),I=1,5)

```

```

WRITE (22,8050) XTIME

```

```

8050 FORMAT (F15.5)

```

```

83 STOP

```

```

END

```